

Shortest networks in Banach spaces:

does existence of a shortest network for every three-point subset of X
imply existence of a shortest network for every finite subset of X ?

Problem stated by Natalia Strelkova

Networks in Banach spaces. Here a *network* in a Banach space is just a finite set of segments. The natural graph structure on a network is defined in the following way. The *edges* are the segments, the *vertices* are the endpoints of the segments. If two segments share an endpoint then the two corresponding edges share the corresponding vertex. A network is called *connected* if the corresponding graph is connected. The *length* $\ell(N)$ of a network N is the sum of the lengths of all its edges (segments), where the length of a segment $[a, b]$ is by definition $\|a - b\|$.

Now let X be a Banach space and fix a finite set $A = \{a_1, \dots, a_n\} \subset X$. We say that a network N *connects* A if N is connected and all points from A are vertices of N . Note that N may also have vertices that are not in A ; we call these vertices *additional*.

Shortest networks. Consider the infimum of length functional $\text{smt}(A) = \inf_N \ell(N)$ over the set of networks that connect A . The general question is whether this infimum is achieved, i.e. whether there exists a *shortest network* — a network that connects A and is shorter than any other network that connects A . It is not difficult to prove that if X is reflexive then for any $A \subset X$ there exists a shortest network. The non-reflexive case is more complicated, and the shortest network may not exist (see below).

Shortest networks with one additional point, or $r_1(A)$. Consider $r_1(A) = \inf_{x \in X} \sum_{i=1}^n \|x - a_i\|$. In fact this is minimizing network length not over the set of all networks connecting A but over a specific subset — the networks with one additional point x and the set of edges $\{[a_i, x]\}_{i=1}^n$. Is the infimum $r_1(A)$ achieved? If X is reflexive then the answer is yes. But in general this is not true, see [1], [2].

Connection between the two existence problems. Note that if A consists of three points then the problem of finding the shortest network is exactly the problem of finding an additional point x at which $r_1(A)$ is achieved.

In [3] it was shown that there exists a Banach space X and for any $n \geq 3$ a set A_n of n points in X such that there

is no shortest network connecting A . The idea of the proof is to take the three-point set A_3 from [2] such that $r_1(A_3)$ is not achieved and prove that for every set A_n that is sufficiently close to A_3 in Hausdorff metric there is no shortest network connecting A_n . (Recall that ε -close in Hausdorff metric means that for any $x \in A_3$ there exists a point $y \in A_n$ such that $\|x - y\| < \varepsilon$ and for any $y \in A_n$ there exists a point $x \in A_3$ such that $\|x - y\| < \varepsilon$.)

PROBLEM. Let X be a Banach space. Suppose that for any three-point set $A_3 \subset X$ there exists a shortest network (i.e. $r_1(A_3)$ is achieved). Is it true that for any $A \subset X$ there exists a shortest network?

References

- [1] L. Vesely, "A characterization of reflexivity in the terms of the existence of generalized centers," *Extracta Mathematicae*, 8 : 2-3, 125 – 131 (1993).
- [2] Borodin P.A., "An example of nonexistence of a Steiner point in a Banach space," *Mat. Zametki*, **87** (4), 514–518 (2010). (in Russian) [English transl. in *Math. Notes*, **87** (4), 485–488 (2010).]
- [3] Bednov B.B., Strelkova N.P., "On the existence problem for shortest networks in Banach spaces," *Mat. zametki*, to appear
- [4] V. Kadets, "Under a suitable renorming every nonreflexive Banach space has a finite subset without a Steiner point," *Matematychni Studii*, 36 : 2, 197 – 200 (2011).